

This Page Is Inserted by IFW Operations
and is not a part of the Official Record.

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(19)



Eur päisch s Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 111 226
A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 83111830.2

(22) Date of filing: 25.11.83


(51) Int. Cl.³: C 07 D 209/48
C 07 D 209/76, C 07 D 401/12
C 07 D 403/12, A 61 K 31/40

(30) Priority: 25.11.82 JP 205545/82


(43) Date of publication of application:
20.06.84 Bulletin 84/25(84) Designated Contracting States:
AT BE CH DE FR GB IT LI NL SE(71) Applicant: Eisai Co., Ltd.
6-10, Kolshikawa 4-chome Bunkyo-ku
Tokyo 112(JP)(72) Inventor: Hirose, Noriyasu
2-36-17, Hiyoshi-cho
Kokubunji-shi Tokyo(JP)(72) Inventor: Souda, Shigeru
1271-317, Ooaza-Ushiku Ushiku-machi
Inashiki-Gun Ibaraki Prefecture(JP)(72) Inventor: Miyake, Kazutoshi
1-56-51, Sakae-cho Ushiku-machi
Inashiki-gun Ibaraki Prefecture(JP)(72) Inventor: Kuriyama, Shizuo
1211-19, Nishitsurugaoka Ohi-machi
Iruma-gun Saitama Prefecture(JP)(72) Inventor: Usuki, Kazuyasu
3-24-14, Aoba-cho
Higashimurayama-shi Tokyo(JP)(72) Inventor: Akiyama, Yasuhiro
1-1730-16, Nakasato
Kiyose-shi Tokyo(JP)(72) Inventor: Nagaoka, Naoko
29-4, Higashi-Arai Yatabemachi
Tsukuba-gun Ibaraki Prefecture(JP)(72) Inventor: Kawashima, Hidetoshi
126-30, Komenoitsujita
Toride-shi Ibaraki Prefecture(JP)(74) Representative: Hansen, Bernd, Dr.rer.nat. et al,
Hoffmann, Eitle & Partner Patentanwälte Arabellastrasse
4
D-8000 München 81(DE)

(54) Carboximide derivatives, process for their production and medicines containing same.


(57) A carboximide derivative represented by the general formula:



wherein X is



Z is alkyl, aralkyl, cycloalkyl, cycloalkylalkyl, alkenyl or arylalkenyl, and Y is a substituted or unsubstituted pyridyl, pyrimidyl or phenyl group, and a pharmaceutically acceptable acid addition salt thereof. The carboximide derivative and acid addition salt are useful for treating diabetes and also for depressing the central nervous system. The carboximide derivative may be produced by reacting a sodium alkoxide represented by the general formula:



wherein X and Y have the same meanings as defined above, with Z-Hal wherein Hal is a halogen and Z has the same meaning as defined above.

Carboximide derivatives, process for their production and medicines containing same

This invention relates to novel carboximide derivatives and their pharmaceutically acceptable acid addition salts having excellent medicinal activities, to processes for their production, to medicines containing the same, and to
5 methods for treating diabetes or depressing the central nervous system by the administration thereof.

The carboximide derivatives and their pharmaceutically acceptable acid addition salts are novel compounds. Their chemical structures are considerably different from those of
10 sulfonyl urea agents, biguanide agents and the like which have been widely used to date.

15 The present inventors have unexpectedly found that these novel carboximide derivatives and their pharmaceutically acceptable acid addition salts have hypoglycemic activities, and are thus useful as medicines for treating diabetes.

A further investigation has resulted in a surprising
20 finding that the carboximide derivatives and their pharmaceutically acceptable acid addition salts also have activities to depress the central nervous system.

An object of this invention is to provide novel carboximide derivatives and their pharmaceutically acceptable
25 acid addition salts which are useful as medicines for treating


diabetes, and also as central nervous system depressants.

Another object of this invention is to provide processes for producing such novel carboximide derivatives and their pharmaceutically acceptable acid addition salts.

5 A further object of this invention is to provide novel pharmaceutical compositions useful for treating diabetes, or as central nervous system depressants, which compositions contain the novel carboximide derivatives or their pharmaceutically acceptable acid addition salts as active
10 ingredients.


A still further object of this invention is to provide methods for treating diabetes or depressing the central nervous system by the administration of the novel carboximide derivatives or their pharmaceutically acceptable acid addition
15 salts.

In one aspect of this invention, there is thus provided a carboximide derivative represented by the general formula (I):




wherein X is

25



cycloalkyl, cycloalkylalkyl, alkenyl or arylalkenyl, and Y is a substituted or unsubstituted pyridyl, pyrimidyl or phenyl group, and a pharmaceutically acceptable acid addition salt thereof.

5 In another aspect of this invention, there is provided a process for producing the carboximide derivative or pharmaceutically acceptable acid addition salt thereof, which comprises reacting a sodium alkoxide represented by the formula (II):



15 wherein X and Y have the same meanings as defined above, with a halide represented by the formula (III):



wherein Hal is a halogen atom and Z has the same meaning as defined above, and, where the pharmaceutically acceptable acid addition salt is desired, further reacting the carboximide derivative with the corresponding acid.


20

In a further aspect of this invention, there is provided a pharmaceutical composition for treating diabetes, or depressing the central nervous system, which

25 comprises, as an active ingredient, the carboximide derivative or pharmaceutically acceptable acid addition salt thereof.

In a still further aspect of this invention, there is provided a method for treating diabetes, which method comprises administering to a patient suffering from diabetes an effective amount of the carboximide derivative or pharmaceutically acceptable acid addition salt thereof.

In a still further aspect of this invention, there is provided a method for depressing the central nervous system of a patient, which method comprises administering to the patient an effective amount of the carboximide derivative or pharmaceutically acceptable salt thereof.



in stereoisomeric forms, in other words, as exo, endo, cis and trans forms. All of these stereoisomers are encompassed by the present invention.

For Z, the alkyl groups include, for example, those containing 1 - 16 carbon atoms, especially straight-chain alkyl containing 1 - 10 carbon atoms; the aralkyl groups include, for example, those where the aryl group is phenyl and the alkyl group contains 1 - 4 carbon atoms, and the aryl group may be substituted, for example, by one or two halogen atoms; the

cycloalkyl groups include, for example, those containing 3 - 7 carbon atoms; the cycloalkylalkyl groups include, for example, those where the cycloalkyl group contains 3 - 7 carbon atoms and the alkyl group contains 1 - 4 carbon atoms; the alkenyl groups include, for example, those containing 2 - 10 carbon atoms; and the arylalkenyl groups include, for example, those where the aryl group is phenyl and the alkenyl group contains 2 - 4 carbon atoms.

For Y, each of the pyridyl, pyrimidyl and phenyl groups may be unsubstituted or substituted. The substituents include, for example, alkyl of 1 - 4 carbon atoms, halogen and trifluoromethyl. There may be, for example, 1 or 2 such substituents.

As mentioned above, a carboximide derivative of the formula (I) may be readily converted to a pharmaceutically acceptable acid addition salt by reacting the derivative with an inorganic or organic acid. As illustrative acids useful for the production of such salts, there may be mentioned inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid and the like, as well as organic acids such as maleic acid, fumaric acid, succinic acid, acetic acid, malonic acid, citric acid, oxalic acid, benzoic acid and the like.

When producing a carboximide derivative of the formula (I) in accordance with the above-described process of this invention, it is acceptable to use, for example, dioxane,

dimethylformamide, dimethylsulfoxide, sulforan or the like as a reaction solvent.

The reaction between compounds (II) and (III) may be carried out under any conditions which are effective to produce the desired compound (I). For example, the reaction may be carried out between 1 - 2 moles of (III) per mole of (II) at a temperature of from room temperature to the boiling point of the solvent until (I) is formed, more particularly, 50 - 100°C for 2 - 12 hours.

10 The compounds according to this invention have excellent glucose tolerance improving activities and are thus useful as medicines for treating diabetes.

In the case of medicines for treating diabetes, long-term continuous administration is indispensable due to the nature of the illness. The compounds according to this invention have low toxicities. Here again, the present invention is believed to be important.

When a compound according to this invention is used as a medicine for treating diabetes or as a central nervous system depressant, it is administered by the oral or parenteral route (i.e., intramuscularly, subcutaneously or intravenously, or as suppositories, or by another suitable administration technique). Although its dosage may vary depending on the severity of the condition of the patient, it generally ranges from 20 mg to 1,000 mg, preferably from 50 mg to 250 mg, per day for a human adult.

In order to form the compounds according to this invention into suitable dosage forms, they are made into such forms as tablets, granules, powders, capsules, injectable solutions, suppositories, etc. in accordance with routine techniques employed in this field.

More specifically, for preparing a solid preparation for oral administration, the active ingredient is mixed with an excipient and, if necessary, further additives such as a binder, disintegrator, lubricant, coloring agent, flavoring agent and the like, and then formed into tablets, coated tablets, granules, powders, capsules, etc. by methods known *per se* in the art.

Examples of the excipients include milk sugar, corn starch, white sugar, glucose, sorbitol, microcrystalline cellulose, etc.. On the other hand, illustrative binders include polyvinyl alcohol, polyvinyl ether, ethylcellulose, methylcellulose, gum arabic, tragacanth gum, gelatin, shellac, hydroxypropylcellulose, hydroxypropylstarch, polyvinyl pyrrolidone and the like. As illustrative disintegrators, there may be mentioned starch, agar, gelatin powder, microcrystalline cellulose, calcium carbonate, sodium hydrogencarbonate, calcium citrate, dextrin, pectin, etc. Lubricants include magnesium stearate, talc, polyethylene glycol, silica, hardened vegetable oil, etc.. Illustrative coloring agents include those permitted for incorporation in medicines. As flavoring agents, use may be made of, for example, cocoa powder, menthol, aromatic acids,

mint oil, camphol, cinnamon powder, etc.. These tablets, granules and the like may of course be suitably coated with sugar, gelatin or the like.

For preparing an injectable solution, the active ingredient is mixed with a pH-adjusting agent, buffer, stabilizer, preservative, etc. and then formed into liquid preparations for subcutaneous, intramuscular or intravenous injection by a method known *per se* in the art.

Examples of the present invention are set forth below in order to describe this invention in more detail. However, the present invention is by no means limited to these examples.

EXAMPLE 1:

15 N-[2-Ethoxy-3-{4-(2-pyridyl)piperazin-1-yl}propyl]-
endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide
dimalate

55% Sodium hydride (0.8 g) was suspended in anhydrous dioxane (50 ml), to which was added dropwise, under nitrogen gas stream and at room temperature, a solution which had been obtained by dissolving N-[2-hydroxy-3-{4-(2-pyridyl)piperazin-1-yl}propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide (5.7 g) in dioxane. The resulting mixture was heated to 60°C and stirred for 1 hour. Thereafter, ethyl iodide (3.0 g) was added dropwis and th thus formed mixture was heated to 75°C and stirred for 5 hours. After cooling the reaction mixture,

25

ice water was added to the reaction mixture. It was extracted with chloroform. The extract was washed with water and then dried with anhydrous magnesium sulfate. It was thereafter filtered and the filtrate was concentrated. The residue was subjected to silica gel chromatography (developer: a 98:2 mixture of chloroform and ethanol) to obtain N-[2-ethoxy-3-{4-(2-pyridyl)piperazin-1-yl}propyl]-endo-cis-bicyclo[2.2.1]-hept-5-ene-2,3-dicarboximide (yield: 49%). It was recrystallized from a mixed solvent of isopropyl ether and n-hexane. Its melting point was 99 - 102°C. A portion (1.64 g) of the thus obtained dicarboximide derivative was then dissolved in ethyl acetate, followed by addition of maleic acid (0.93 g) dissolved in a mixed solvent of ethyl acetate and methanol. The resulting mixture was heated. By allowing the reaction mixture to cool, the desired compound was caused to precipitate. It was collected by filtration (yield: 2.45 g).

Melting point (°C): 136 - 138

Elemental Analysis

for $C_{23}H_{30}N_4O_3 \cdot 2C_4H_4O_4$

		<u>C</u>	<u>H</u>	<u>N</u>
20	Calculated (%)	57.93	5.97	8.72
	Found (%)	57.85	5.95	8.74

EXAMPLE 2:

N-[2-n-Butoxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-
endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide
dimaleate

5 55% sodium hydride (1.05 g) was suspended in anhydrous dioxane (60 ml), to which a solution of N-[2-hydroxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl-endo-cis-bicyclo[2.2.1]hept-5-en-2,3-dicarboximide (7.64 g) in dioxane was added dropwise under nitrogen gas stream and at room temperature. The resulting
10 mixture was heated to 65°C and stirred for 1 hour. n-Butyl bromide (3.2 g) was added dropwise at 40°C and the resulting mixture was agitated for 6 hours at 75 - 80°C. After cooling the reaction mixture, ice water was added thereto and it was then extracted with ether. The extract was washed with water
15 and then dried with anhydrous magnesium sulfate. It was filtered and the filtrate was concentrated. Mineral oil contained in the residue, derived from sodium hydride, was washed off with n-hexane. The yield of the thus obtained N-[2-n-butoxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide was 3.7 g (yield: 42%). It was then
20 taken up in ethyl acetate, followed by addition of maleic acid (2.0 g) dissolved in a mixed solvent of ethyl acetate and methanol. The resulting mixture was then heated. The resulting reaction mixture was allowed to cool and the desired compound
25 was hence caused to precipitate. The desired compound was

collected by filtration, and had the following physical properties (yield: 4.9 g).

Melting point (°C): 126 - 129

Elemental Analysis: for $C_{25}H_{34}N_4O_3 \cdot 2C_4H_4O_4$:

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	59.08	6.32	8.35
Found (%)	58.81	6.32	8.44

EXAMPLE 3:

N-[2-Cyclohexylmethoxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide dimaleate

55% Sodium hydride (0.85 g) was suspended in anhydrous dioxane (50 ml), followed by dropwise addition of N-[2-hydroxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide (5.7 g) dissolved in dioxane (60 ml) under nitrogen gas stream and at room temperature. The resulting mixture was heated to 65 - 70°C and stirred for 1 hour. Cyclohexylmethyl bromide (4.0 g) was added dropwise at 40°C and the thus formed mixture was stirred at 80 - 85°C for 10 hours. After cooling the reaction mixture, ice water was added to the mixture and the resulting mixture was extracted with ether. The extract was washed with water and then dried with anhydrous magnesium sulfate. It was filtered and the filtrate was concentrated. The residue was subjected

to silica gel chromatography (developer: a 98:2 mixture of chloroform and ethanol) to obtain 3.9 g of N-[2-cyclohexylmethoxy-3-(4-(2-pyridyl)-piperazin-1-yl)propyl]-endo-cis-bicyclo [2.2.1]hept-5-ene-2,3-dicarboximide [Rf value: 0.5 (developer: a 5 96:4 mixture of chloroform and ethanol); yield: 54%]. The reaction product was then dissolved in ethyl acetate, followed by addition of a solution which had been obtained by dissolving maleic acid (1.9 g) in a mixed solvent of ethyl acetate and methanol. The resulting mixture was heated. It 10 was then allowed to cool to cause the desired compound to precipitate. The precipitate was collected by filtration to obtain the desired compound having the following physical properties (yield: 4.5 g).

Melting point (°C): 137 - 140

15-

Elemental Analysis

for $C_{28}H_{38}N_4O_3 \cdot 2C_4H_4O_4$:

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	60.82	6.53	7.88
Found (%)	60.82	6.49	7.83

20

EXAMPLE 4:

N-[2-Allyloxy-3-(4-(2,4-dimethylphenyl)piperazin-1-yl)propyl]-phthalimide oxalate

55% Sodium hydride (0.65 g) was suspended in anhydrous 25 dioxane (30 ml), followed by dropwise addition of N-[2-hydroxy-3-(4-(2,4-dimethylphenyl)piperazin-1-yl)propyl]-phthalimide

(4.7 g) dissolved in dioxane (50 ml) under nitrogen gas-stream and at room temperature. The resulting mixture was heated to 60°C and stirred for 1 hour. Then, allyl bromide (1.8 g) was dropped slowly into the mixture at 40°C and the thus formed mixture was stirred for 3 hours over a water bath at 75°C. The reaction mixture was thereafter cooled, ice water was added thereto, and it was then extracted with ether. After washing the extract twice with water, it was dried with anhydrous magnesium sulfate. The resulting mixture was filtered and the filtrate was concentrated. After washing out the mineral oil, derived from sodium hydride, with n-hexane, the residue was dissolved in ethyl acetate, followed by addition of a solution which had been obtained by dissolving oxalic acid (1.2 g) in ethyl acetate. After heating the resulting mixture, a small amount of IPE (isopropyl ether) was added. The thus obtained mixture was allowed to cool. Precipitated crystals were collected by filtration to obtain the desired compound (4.3 g; yield 69%).

Melting point (°C): 146 - 149

Elemental Analysis for $C_{26}H_{31}N_3O_3 \cdot C_2H_2O_4$:

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	64.22	6.37	8.03
Found (%)	64.15	6.12	7.90

EXAMPLE 5:

N-(2-Ethoxy-3-(4-(3-chlorophenyl)piperazin-1-yl)-propyl)-endo-cis-bicyclo[2.2.1]heptane-2,3-dicarboximide oxalate

5 N-(2-Ethoxy-3-(4-(3-chlorophenyl)piperazin-1-yl)-propyl)-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide (1.7 g) was dissolved in a mixed solvent of ethanol (30 ml) and ethyl acetate (10 ml), and then catalytically reduced using 10% palladium-carbon as a catalyst. After confirming
10 by T.L.C. that the raw material had been completely used up, the catalyst was separated by filtration and the filtrate was concentrated. The residue was dissolved in ethyl acetate, followed by addition of a solution containing oxalic acid (500 mg) in ethyl acetate. The resulting mixture was heated
15 to dissolve solid materials completely. A small amount of isopropyl ether was added to the reaction mixture, which was then finally cooled to obtain the desired compound (1.9 g; yield: 74%).

Melting point (°C) 158 - 160

20	Elemental Analysis	for $C_{24}H_{32}N_3O_3Cl \cdot C_2H_2O_4$:		
		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	58.25	6.41	7.84
	Found (%)	58.28	6.32	7.92

25 Following the procedure of Example 1, the following compound was obtained:

N-[2-Ethoxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]phthalimide dimaleate

Melting point (°C): 135 - 137

Elemental Analysis for $C_{22}H_{26}N_4O_3 \cdot 2C_4H_4O_4$:

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	57.49	5.48	8.94
Found (%)	57.25	5.40	8.96

Repeating the procedure of Example 2, the following compounds were obtained.

N-[2-Benzoyloxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide dimaleate

Melting point (°C): 114 - 117

Elemental Analysis for $C_{28}H_{32}N_4O_3 \cdot 2C_4H_4O_4$:

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	61.35	5.73	7.95
Found (%)	61.07	5.64	7.92

N-[2-Allyloxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.2]oct-5-ene-2,3-dicarboximide dimaleate

Melting point (°C) 114 - 117

Elemental Analysis for $C_{25}H_{32}N_4O_3 \cdot 2C_4H_4O_4$:

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	59.26	6.04	8.38
Found (%)	59.02	6.02	8.20

N-[2-(4-Chlorobenzoyloxy)-3-{4-(2-pyrimidyl)piperazin-1-yl}-propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide maleate

Melting point (°C): 162 - 164

5 Elemental Analysis for $C_{27}H_{30}ClN_5O_3 \cdot C_4H_4O_4$:

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	59.65	5.50	11.22
Found (%)	59.74	5.27	11.14

10

Following the procedure of Example 3, the following compounds were obtained.

N-[2-(n-Octyloxy)-3-{4-(2-pyridyl)piperazin-1-yl}propyl]-cis-cyclohex-4-ene-1,2-dicarboximide 1.7 oxalate

Melting point (°C): 109 - 112

15 Elemental Analysis for $C_{29}H_{43}N_3O_3 \cdot 1.7C_2H_2O_4$:

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	61.30	7.38	6.62
Found (%)	61.02	7.53	6.65

20 N-[2-n-Butoxy-3-{4-(3-chlorophenyl)piperazin-1-yl}propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide oxalate

Melting point (°C): 122 - 125

Elemental Analysis for $C_{26}H_{34}ClN_3O_3 \cdot C_2H_2O_4$:

	<u>C</u>	<u>H</u>	<u>N</u>
25 Calculated (%)	59.82	6.47	7.48
Found (%)	59.62	6.58	7.35

N-[2-Ethoxy-3-(4-(3-chlorophenyl)piperazin-1-yl)propyl]-
endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide oxalate

Melting point (°C): 159 - 161

Elemental Analysis for $C_{24}H_{30}ClN_3O_3 \cdot C_2H_2O_4$:

5		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	58.47	6.05	7.87
	Found (%)	58.27	6.04	7.83

N-[2-n-Decyloxy-3-(4-(3-chlorophenyl)piperazin-1-yl)propyl]-
endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide oxalate

Melting point (°C): 119 - 122

Elemental Analysis for $C_{32}H_{46}ClN_3O_3 \cdot C_2H_2O_4$:

10		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	63.18	7.50	6.50
15	Found (%)	63.03	7.37	6.51

N-[2-n-Butoxy-3-(4-(3-chlorophenyl)piperazin-1-yl)propyl]-
cis-cyclohex-4-ene-1,2-dicarboximide 1.1 oxalate

Melting point (°C): 116 - 119

Elemental Analysis for $C_{25}H_{34}ClN_3O_3 \cdot 1.1C_2H_2O_4$:

20		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	58.21	6.52	7.51
	Found (%)	58.09	6.69	7.36

25 .

N-[2-n-Butoxy-3-(4-(4-chlorophenyl)piperazin-1-yl)propyl]-
endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide oxalate

Melting point (°C): 152 - 155

Elemental Analysis for $C_{26}H_{34}ClN_3O_3 \cdot C_2H_2O_4$:

5		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	59.82	6.47	7.48
	Found (%)	59.60	6.61	7.46

10 N-[2-n-Butoxy-3-(4-(2-chlorophenyl)piperazin-1-yl)-propyl]-
endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide dioxalate

Melting point (°C): 109 - 112

Elemental Analysis for $C_{26}H_{34}ClN_3O_3 \cdot 2C_2H_2O_4$:

		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	55.27	5.89	6.45
15	Found (%)	55.49	6.10	6.67

The procedure of Example 4 was repeated to obtain

the following compounds.

N-[2-Allyloxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]phthalimide
1.2 oxalate

20 Melting point (°C): 130 - 133

Elemental Analysis for $C_{25}H_{29}N_3O_4 \cdot 1.2C_2H_2O_4$:

		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	60.09	5.80	7.73
25	Found (%)	60.09	5.80	7.63

N-[2-Benzoyloxy-3-(4-(3-chlorophenyl)piperazin-1-yl)propyl]-
endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide oxalate

Melting point (°C): 168 - 170

Elemental Analysis for $C_{29}H_{32}ClN_3O_3 \cdot C_2H_2O_4$:

5

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	62.45	5.76	7.05
Found (%)	62.71	5.81	6.96

N-[2-Cinnamyloxy-3-(4-(α,α,α -trifluoro-3-tolyl)piperazin-1-
10 yl)-propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide
oxalate

Melting point (°C): 164 - 166

Elemental Analysis for $C_{32}H_{34}F_3N_3O_3 \cdot C_2H_2O_4$:

15

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	62.27	5.54	6.41
Found (%)	62.38	5.57	6.41

Following the procedure of Example 5, the following
compounds were obtained.

20

N-[2-n-Octyloxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-
cyclohexane-1,2-dicarboximide 1.7 oxalate

Melting point (°C): 120 - 123

Elemental Analysis for $C_{29}H_{45}N_3O_3 \cdot 1.7C_2H_2O_4$:

25

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	61.10	7.68	6.60
Found (%)	61.00	7.65	6.63

N-[2-n-Butoxy-3-{4-(3-chlorophenyl)piperazin-1-yl}propyl]-
endo-cis-bicyclo[2.2.1]heptane-2,3-dicarboximide oxalate

Melting point (°C): 123 - 125

Elemental Analysis for $C_{26}H_{36}ClN_3O_3 \cdot C_2H_2O_4$:

5		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	59.61	6.80	7.45
	Found (%)	59.49	6.77	7.48

N-[2-n-Decyloxy-3-{4-(3-chlorophenyl)piperazin-1-yl}propyl]-
endo-cis-bicyclo[2.2.1]heptane-2,3-dicarboximide oxalate

Melting point (°C): 120 - 123

Elemental Analysis for $C_{32}H_{48}ClN_3O_3 \cdot C_2H_2O_4$:

15		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	62.98	7.79	6.48
	Found (%)	62.93	7.66	6.57

N-[2-n-Butoxy-3-{4-(4-chlorophenyl)piperazin-1-yl}propyl]-
endo-cis-bicyclo[2.2.1]heptane-2,3-dicarboximide oxalate

Melting point (°C): 152 - 155

Elemental Analysis for $C_{26}H_{36}ClN_3O_3 \cdot C_2H_2O_4$:

20		<u>C</u>	<u>H</u>	<u>N</u>
	Calculated (%)	59.61	6.80	7.45
	Found (%)	59.22	7.03	7.34

25

N-[2-n-Butoxy-3-{4-(2-chlorophenyl)piperazin-1-yl}propyl]-
endo-cis-bicyclo[2.2.1]heptane-2,3-dicarboximide 1.7 oxalate

Melting point (°C): 108 - 111

Elemental Analysis for $C_{26}H_{36}ClN_3O_3 \cdot 1.7C_2H_2O_4$:

5

	<u>C</u>	<u>H</u>	<u>N</u>
Calculated (%)	56.29	6.35	6.70
Found (%)	56.03	6.40	6.79

Next, exemplary preparations containing representa-
 10 tive compounds of this invention will be described.

Preparation Example 1: Tablets

	N-[2-Ethoxy-3-{4-(2-pyridyl)piperazin- 1-yl}propyl]phthalimide dimaleate	50 g
15	Corn starch	10 g
	Milk sugar	65 g
	Calcium carboxymethylcellulose	10 g
	Polyvinyl pyrrolidone	5 g
	Talc	10 g
20	Microcrystalline cellulose	50 g

In accordance with methods known *per se* in the
 art, all the above ingredients were mixed, granulated and
 then press-formed into tablets, each of 200 mg.

25

Preparation Example 2: Capsules

	N-[2-n-Butoxy-3-{4-(3-chlorophenyl)-	50 g
	piperazin-1-yl}propyl]-cis-cyclohex-4-	
	ene-1,2-dicarboximide 1.1 oxalate	
5	Milk sugar	45 g
	Corn starch	5 g

The above composition was prepared into capsules,
each containing 100 mg, in a manner known *per se* in the art.

Having now fully described the invention, it will
10 be apparent to one of ordinary skill in the art that many
changes and modifications can be made thereto without departing
from the spirit or scope of the invention as set forth herein.


15

20


25

CLAIMS:

1. A compound selected from the group consisting of a carboximide of the formula:



wherein



15 Z is alkyl, aralkyl, cycloalkyl, cycloalkylalkyl, alkenyl or arylalkenyl, and

Y is a substituted or unsubstituted pyridyl, pyrimidyl or phenyl group, and


a pharmaceutically acceptable acid addition salt thereof.

20

2. A compound according to claim 1, wherein X is



or



25

3. A compound according to claim 1, wherein Y is 2-pyridyl.
4. A compound according to claim 1, wherein Y is 2-chlorophenyl, 3-chlorophenyl or 4-chlorophenyl.
- 5
5. A compound according to claim 1, wherein Y is 2,4-dimethylphenyl.
6. A compound according to claim 1, wherein Y is α, α, α -tri-
10 fluoro-3-tolyl.
7. A compound according to claim 1, wherein Z is a straight-chain alkyl group containing 1 - 10 carbon atoms.
- 15 8. A compound according to claim 7, wherein the straight-chain alkyl group is ethyl, n-butyl, n-octyl or n-decyl.
9. A compound according to claim 1, wherein Z is substituted or unsubstituted benzyl.
- 20
10. A compound according to claim 1, wherein Z is allyl.
11. A compound according to claim 1, wherein Z is cinnamyl.
- 25 12. A pharmaceutically acceptable acid addition salt according to claim 1, wherein the salt is a hydrochloride, hydrobromide, sulfate, maleate, fumarate, succinate, acetate, malonate, citrate, oxalate or benzoate.

13. A compound according to claim 1, which is N-[2-ethoxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]-hept-5-ene-2,3-dicarboximide or the dimaleate thereof.

5 14. A compound according to claim 1, which is N-[2-n-butoxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]-hept-5-ene-2,3-dicarboximide or the dimaleate thereof.

15. A compound according to claim 1, which is N-[2-cyclohexyl-10 methoxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-endo-cis-bicyclo-[2.2.1]hept-5-ene-2,3-dicarboximide or the dimaleate thereof.

16. A compound according to claim 1, which is N-[2-allyloxy-3-(4-(2,4-dimethylphenyl)piperazin-1-yl)propyl]-phthalimide
15 or the oxalate thereof.

17. A compound according to claim 1, which is N-[2-ethoxy-3-(4-(3-chlorophenyl)piperazin-1-yl)propyl]-endo-cis-bicyclo-[2.2.1]heptane-2,3-dicarboximide or the oxalate thereof.
20

18. A compound according to claim 1, which is N-[2-ethoxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-phthalimide or the dimaleate thereof.

25 19. A compound according to claim 1, which is N-[2-benzyloxy-3-(4-(2-pyridyl)piperazin-1-yl)propyl]-endo-cis-bicyclo-[2.2.1]hept-5-ene-2,3-dicarboximide or the dimaleate thereof.

20. A compound according to claim 1, which is N-[2-allyloxy-3-{4-(2-pyridyl)piperazin-1-yl}propyl]-endo-cis-bicyclo[2.2.2]-oct-5-ene-2,3-dicarboximide or the dimaleate thereof.
- 5 21. A compound according to claim 1, which is N-[2-(4-chlorobenzyloxy)-3-{4-(2-pyrimidyl)piperazin-1-yl}propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide or the maleate thereof.
- 10 22. A compound according to claim 1, which is N-[2-(n-octyloxy)-3-{4-(2-pyridyl)piperazin-1-yl}propyl]-cis-cyclohex-4-ene-1,2-dicarboximide or the 1.7 oxalate thereof.
- 15 23. A compound according to claim 1, which is N-[2-n-butoxy-3-{4-(3-chlorophenyl)piperazin-1-yl}propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide or the oxalate thereof.
24. A compound according to claim 1, which is N-[2-ethoxy-3-{4-(3-chlorophenyl)piperazin-1-yl}propyl]-endo-cis-bicyclo-
20 [2.2.1]hept-5-ene-2,3-dicarboximide or the oxalate thereof.
25. A compound according to claim 1, which is N-[2-n-decyloxy-3-{4-(3-chlorophenyl)piperazin-1-yl}propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide or the oxalate thereof.

26. A compound according to claim 1, which is N-[2-n-butoxy-3-(4-(3-chlorophenyl)piperazin-1-yl)propyl]-cis-cyclohex-4-ene-1,2-dicarboximide or the 1.1 oxalate thereof.

5 27. A compound according to claim 1, which is N-[2-n-butoxy-3-(4-(4-chlorophenyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide or the oxalate thereof.

28. A compound according to claim 1, which is N-[2-n-butoxy-3-10 -(4-(2-chlorophenyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]-hept-5-ene-2,3-dicarboximide or the dioxalate thereof.


29. A compound according to claim 1, which is N-[2-allyloxy-3-15 -(4-(2-pyridyl)piperazin-1-yl)propyl]phthalimide or the 1.2 oxalate thereof.

30. A compound according to claim 1, which is N-[2-benzyloxy-20 -3-(4-(3-chlorophenyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide or the oxalate thereof.


31. A compound according to claim 1, which is N-[2-cinnamyloxy-3-(4-(α,α,α -trifluoro-3-tolyl)piperazin-1-yl)propyl]-endo-cis-bicyclo[2.2.1]hept-5-ene-2,3-dicarboximide or the oxalate thereof.

25

32. A compound according to claim 1, which is N-[2-n-octyloxy-3-{4-(2-pyridyl)piperazin-1-yl}propyl]-cyclohexane-1,2-dicarboximide or the 1.7 oxalate thereof.
- 5 33. A compound according to claim 1, which is N-[2-n-butoxy-3-{4-(3-chlorophenyl)piperazin-1-yl}propyl]-endo-cis-bicyclo-[2.2.1]heptane-2,3-dicarboximide or the oxalate thereof.
34. A compound according to claim 1, which is N-[2-n-decyloxy-10 -3-{4-(3-chlorophenyl)piperazin-1-yl}propyl]-endo-cis-bicyclo-[2.2.1]heptane-2,3-dicarboximide or the oxalate thereof.
35. A compound according to claim 1, which is N-[2-n-butoxy-15 -3-{4-(4-chlorophenyl)piperazin-1-yl}propyl]-endo-cis-bicyclo-[2.2.1]heptane-2,3-dicarboximide or the oxalate thereof.
36. A compound according to claim 1, which is N-[2-n-butoxy-20 -3-{4-(2-chlorophenyl)piperazin-1-yl}propyl]-endo-cis-bicyclo-[2.2.1]heptane-2,3-dicarboximide or the 1.7 oxalate thereof.
37. A process for producing a carboximide of the formula



wherein




Z is alkyl, aralkyl, cycloalkyl, cycloalkylalkyl, alkenyl or arylalkenyl, and


Y is a substituted or unsubstituted pyridyl, pyrimidyl or phenyl group,

or a pharmaceutically acceptable acid addition salt thereof,

which comprises reacting a sodium alkoxide of the formula:



wherein X and Y have the same meanings as defined above, with a halide of the formula:











wherein Hal is a halogen atom and Z has the same meaning as defined above, and, where the pharmaceutically acceptable acid addition salt is desired, further reacting the resultant carb-
boximide with an acid.













38. A process according to claim 37, wherein the reaction is carried out in a solvent selected from the group consisting of dioxane, dimethylformamide, dimethylsulfoxide and sulforan.




5 39. A process according to claim 37, wherein the acid is selected from the group consisting of hydrochloric acid, hydrobromic acid, sulfuric acid, maleic acid, fumaric acid, succinic acid, acetic acid, malonic acid, citric acid, oxalic acid and benzoic acid.

10


40. A process according to claim 37, wherein X, Y, Z and Hal are selected from the following combinations:

	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>Hal</u>
15		2-pyridyl	ethyl	I
		2-pyridyl	n-butyl	Br
		2-pyridyl	cyclohexylmethyl	Br
20		2,4-dimethylphenyl	allyl	Br
		2-pyridyl	ethyl	I
		2-pyridyl	benzyl	Br
		2-pyridyl	allyl	Br
25		2-pyrimidyl	4-chlorobenzyl	Br


	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>Hal</u>
5		2-pyridyl	n-octyl	Br
		3-chlorophenyl	n-butyl	Br
		3-chlorophenyl	ethyl	Br
		3-chlorophenyl	n-decyl	Br
		3-chlorophenyl	n-butyl	Br
10				
		4-chlorophenyl	n-butyl	Br
		2-chlorophenyl	n-butyl	Br
		2-pyridyl	allyl	Br
		3-chlorophenyl	benzyl	Br
15		α, α, α -trifluoro-3-tolyl	cinnamyl	Br
		2-pyridyl	n-octyl	Br
		3-chlorophenyl	n-butyl	Br
20				
25				

<u>X</u>	<u>Y</u>	<u>Z</u>	<u>Hal</u>
	3-chlorophenyl	n-decyl	Br
	4-chlorophenyl	n-butyl	Br
	2-chlorophenyl	n-butyl	Br

41. A pharmaceutical composition for treating diabetes which comprises, as an active ingredient, a carboximide of the formula:



wherein




Z is alkyl, aralkyl, cycloalkyl, cycloalkylalkyl, alkenyl or arylalkenyl, and


Y is a substituted or unsubstituted pyridyl, pyrimidyl or phenyl group, or a pharmaceutically acceptable acid addition salt thereof, and

a pharmaceutically acceptable carrier therefor.

42. A central nervous system depressant composition which comprises, as an active ingredient, a carboximide of the formula:



wherein



Z is alkyl, aralkyl, cycloalkyl, cycloalkylalkyl, alkenyl or alylalkenyl, and

Y is a substituted or unsubstituted pyridyl, pyrimidyl or phenyl group,

20 or a pharmaceutically acceptable acid addition salt thereof, and

a pharmaceutically acceptable carrier therefor.



European Patent
Office

EUROPEAN SEARCH REPORT

0111226

Application number

EP 83 11 1830

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
Y	US-A-2 672 460 (E.A. CONROY) * Whole document *	1-12	C 07 D 209/48 C 07 D 209/76 C 07 D 401/12 C 07 D 403/12 A 61 K 31/40
Y	--- US-A-2 343 198 (M.B. MOORE) * Whole document *	1-12	
P,X	--- FR-A-2 506 771 (ESAI CO. LTD.) * Pages 1-5; claims *	1-42	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 7)
			C 07 D 209/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 07-03-1984	Examiner MAISONNEUVE J.A.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO Form 1503 03/82

THIS PAGE BLANK (USPTO)